



Rte Ancienne Papeterie
PO Box 327,
CH-1723 Marly 1 (Switzerland)

Fax: +41 91 825 76 08
info@swant.com
www.swant.com

Product Description

Monoclonal anti Na⁺-Ca²⁺-exchanger

Product: Monoclonal anti-Na⁺-Ca²⁺-exchanger

Code No: R3F1

Lot no: #mr05

Form: Lyophilized, concentrated supernatant.

IgG Subtype: IgG1

Quantity: 200 µl

Reconstitution: with 200 µl of bidistilled water.

Description

The monoclonal antibody against the canine Na⁺-Ca²⁺-exchanger, recognizes two neighboring but non-overlapping sequences of the hydrophilic region of the exchanger, connecting the putative transmembrane segments 5 and 6. The antibody was evaluated for specificity and potency by immunoblotting, and immunoprecipitation (see reference). It recognizes virtually only NCX1 in monkey, horse, dog, rat, bovine and mouse tissue. At higher concentrations it reacts faintly with NCX2, whereas NCX3 is not recognized. Both splice variants with exon A (heart, nervous system) and exon B (kidney, glia) are recognized.

Working dilutions

Immunohistochemistry: 1: 500; 1: 1'000

Immunoblots and immunoprecipitation: 1: 500; 1: 1'000 (use milk for blocking). A strong 70 kDa band is found in addition to the 116 kDa of the integral exchanger. Often the HMW band is a double band 116/ 112 kDa band.

The titer was determined using the indirect-peroxidase method. We recommend that the optimal dilutions be determined by titration experiments.

Storage

After reconstitution freeze in small aliquots (e.g. 1 µl) and keep at - 80°C (or at least - 20°C). For continuous use keep at 4°C (with 0.01% Na-azide). Avoid repeated freezing and thawing.

References

Porzig H., Li Z., Nicoll D.A., Philipson K.D. (1993) Mapping the cardiac sodium-calcium exchanger with monoclonal antibodies. *Am. J. Physiol.* 265:C78 -C756, 1993